

Plate Tectonics: too weak to build mountains

For this discussion, the assumptions and ideas of plate tectonics are used unchallenged to show their problems regarding mountain building (orogeny). Quotes are from professional journals.

What drives the plates?

Study of the motions of plates is called kinematics, while study of the driving forces is called dynamics. "A key to the simplicity of plate tectonics is that the strength of lithospheric plates enables the analysis of their kinematics to be isolated and treated separately from the dynamic processes controlling plate motions; relative velocities of plates can be analyzed without reference to the forces that give rise to them"²⁸.

Around the end of the first decade of dominance by plate tectonics, in 1975, the situation was described this way: "In recent years, the kinematics of continental drift and sea-floor spreading have been successfully described by the theory of plate tectonics. However, rather little is known about the driving mechanisms of plate tectonics, although various types of forces have been suggested"¹². Seven years later, in 1982, the assessment was: "At the present time the geometry of plate movements is largely understood, but the driving mechanism of plate tectonics remains elusive"¹. By 1995 we find that: "In spite of all the mysteries this picture of moving tectonic plates has solved, it has a central, unsolved mystery of its own: What drives the plates in the first place?'[That] has got to be one of the more fundamental problems in plate tectonics,' notes geodynamicist Richard O'Connell of Harvard University. 'It's interesting it has stayed around so long' "²⁰. In 2002 it could be said that: "Although the concept of plates moving on Earth's surface is universally accepted, it is less clear which forces cause that motion. Understanding the mechanism of plate tectonics is one of the most important problems in the geosciences"⁶. A 2004 paper noted that "considerable debate remains about the driving forces of the tectonic plates and their relative contribution"³³. "Alfred Wegener's theory of continental drift died in 1926, primarily because no one could suggest an acceptable driving mechanism. In an ironical twist, continental drift (now generalized to plate tectonics) is almost universally accepted, but we still do not understand the driving mechanism in anything other than the most general terms"².

The problem has always been that it is hard to discern what is going on deep in the Earth, motion is almost imperceptibly slow, and different combinations of forces, perhaps varying over time, could apply to particular areas. "When the concepts of convection and plate tectonics were first developing, many thought of mantle convection as a process heated from below, which in turn exerts driving tractions on the base of a relatively stagnant 'crust' (later, 'lithosphere') to cause continental drift. In the early 1970s, more sophisticated understanding of convection led to the opposite view. It was realized that only a fraction of the Earth's heat flow originates in the core, while most results from radioactivity and/or secular cooling of the mantle. Computer models showed that internally heated (and/or surface cooled) systems have no upwelling sheets or plumes and that all concentrated flow originates in the upper cold boundary layer, which stirs the interior as it sinks. Thus it became natural to regard plates of lithosphere as driving themselves and, incidentally, stirring the rest of the mantle"³. Some researchers make the point emphatically: "convection does not drive plates." Upper mantle convection is a product, not a cause, of plate motions¹⁶. Thus the location and orientation of a sinking slab is the best indicator of which way upper mantle flows.

"The advent of plate tectonics made the classical mantle convection hypothesis even more untenable. For instance, the supposition that mid-oceanic ridges are the site of upwelling and trenches are that of sinking of the large scale convective flow cannot be valid, because it is now established that actively spreading, oceanic ridges migrate and often collide with trenches"¹². "Another difficulty is that if this is currently the main mechanism, the major convection cells would have to have about half the width of the large oceans, with a pattern of motion that would have to be more or less constant over very large areas under the lithosphere. This would fail to explain the relative motion of plates with irregularly shaped margins at the Mid-Atlantic ridge and Carlsberg ridge, and the motion of small plates, such as the Caribbean and the Philippine plates"¹⁵.

The available options

Of the possible driving forces, a consensus has developed "that the dominant forces might operate either (1) from the side by 'slab pull' by the subducting plates (slabs) and 'ridge push' from mid-ocean ridges or (2) from below by mantle convection"⁶. "It is a simple matter to ascribe the driving force to gravity causing plates to slide downhill from mid-ocean ridges and pulling them into the asthenosphere at subduction zones, but it is a rare fluid dynamicist who would contend that these processes are understood"²⁸. Ridge push has an additional meaning: the expansion of oceanic crust as it cools and thickens for up to 90 million years²⁶.

"Oceanic plates [are] underlain by the low-viscosity zone (LVZ) that might be 50-200 km thick and in which the coefficient of viscosity is at least an order of magnitude less than that of the mantle in general. Consequently, the possible coupling between a mantle convection cell and the overlying plate [is] seen to be exceedingly improbable. In motoring terms, there [is] a 'slipping clutch' between the engine and the drive shaft"³¹. "The continental and oceanic lithospheres may behave differently, as the LVZ is less well defined, or even non-existent, below continents. Moreover, the continental lithosphere may also have a relatively deep 'keel'. If such a keel exists beneath a continental lithosphere, and plate motion is independent of mantle flow direction, then it is reasonable to infer that the basal force will resist movement in plates containing a large proportion of continental material, more than in plates dominated by oceanic lithosphere. Hence, one would expect plates with a high percentage of continental crust to move relatively slowly. For current major plate geometries this expectation is generally supported"³¹. "Over the past three decades there has been vigorous debate over how thick the continents can be -- that is, the depth to which the rigid crust and upper mantle reach before meeting convecting mantle that can flow and drive tectonic motion"¹⁹. The depths in question are 400 km versus 250 km. Recent work indicates "that continental roots do not extend much beyond a depth of 250 km"¹⁴. The observation that plates with the longest trench boundaries move the fastest points to the importance of slab pull.³¹

Numerical models compare a variety of factors. In some cases, forward basal drag by mantle flow is indicated. "In the most plausible model, this forward drag acts only on continents, while oceanic lithosphere experiences negligible basal shear tractions. Probably the dense descending slabs of oceanic lithosphere not only pull the oceanic plates, but also stir the more viscous lower mantle, and this in turn helps to drive the slower drift of continents"³.

How capable is each option?

In dynamics research, force is described in units of newtons per meter (N/m), pascals (Pa), and bars (b or bar). For clarity, bars will be used here. Estimates of net force generated by each mechanism are

fairly consistent among researchers. For example,

Slab pull: 500 bars³¹, 450 bars ("subduction pull"⁷)

Ridge push: 200 bars²⁶, 250 bars³⁸, 250 bars⁷, 200-300 bars⁴⁰, 200-400 bars³¹


Forward basal drag: 200 bars³⁷, 200 bars³

The force generated by a sinking slab would be much higher, but its net effect is greatly reduced by various forms of resistance. This is especially so because the subduction trench retreats seaward (rollback). A common "misconception is that subducting plates roll over stationary hinges and slide down fixed slots"¹⁶, the way an escalator descends. A closer illustration would be a strip of paint peeling from a ceiling, although there is also a small amount of forward sliding - the net slab pull. There is general consensus for the net slab pull force listed above³¹. Laboratory experiments show that about 70% of the sinking slab's force is used to drive rollback-induced mantle flow; roughly 15 to 30% is used to bend the subducting plate at the trench; and 0 to 8% is used to overcome shear resistance between the slab and the mantle. The experiments indicate only 8 to 12% of the force of the sinking slab pulls the attached surface plate forward. Thus slab pull is about twice as large as the ridge push force³³. We can avoid uncertainties regarding slab pull by considering the case of the South American plate.

South America - a simple example

"The Andes are the world's second largest mountain belt."²¹ "Dynamic analysis of the South American plate is straightforward because of the relative simplicity of the plate's boundaries and the near absence of slab-pull"³². That leaves ridge push and mantle flow coupled to the continent's base as the available forces. "During the last 30 million years, [South America's absolute westward] velocity increased from 2.0 to 2.8 cm/year. South America is currently moving faster relative to the hot spots than at any time in the last 80 million years"³⁶. "The South American margin, despite a geologic history of more than 200 million years of continuous subduction, did not begin to grow high topography until ~35 million years ago." "There is now a consensus that the thick crust and high topography in the Andes essentially reflect... tectonic shortening, quite similar to collisional orogens" (mountains). "Maximum shortening values of 250-300 km are sufficient to thicken the crust of the central Andes to its present state."²¹

"Virtually all major mountain ranges in the world are a consequence of crustal shortening. To form mountain ranges, in general, horizontal forces must be applied to masses of crust and mantle, to lithospheric plates, to drive them together and to cause crustal shortening and crustal thickening"³⁰. "The Andes shouldn't be there. Plate tectonics makes the world's great mountain ranges by slamming two continents together, as Europe collided with Africa to make the Alps or India ran into Asia to make the Himalayas. South America, however, is colliding with nothing more than the floor of the Pacific Ocean, which is slipping beneath the continent into Earth's interior. Such encounters between continent and ocean ordinarily throw up a few volcanoes, not a 7000-kilometer-long wall of mountains"²⁰. A few researchers think South America is colliding with "the viscous mantle rock hundreds of kilometers down under the floor of the Pacific. Like a snub-nosed boat driven too fast for the strength of its hull, the central South American coastline has crumpled under the pressure"²⁰. But what is driving the boat? The forward basal drag from trench suction just keeps the Pacific and Atlantic plates (including South America) from separating.




The stress required for crustal shortening to build mountains has been calculated to be in a range from 1500 to 2500 bars¹⁰ up to 4000 to 6000 bars³¹, inferring the latter "from earthquake data and evaluation of the stresses required to produce specific geological structures". In the case of South America, the combination of ridge push and forward basal drag (by trench suction) could produce only 400 to 600 bars of force, which is clearly insufficient to build the Andes. These forces are already engaged in moving the entire plate westward! A researcher who acknowledged this failure was moved to suggest another mechanism (gravity glide) that had been discarded many years ago³¹ (and is still out of favor). At present, plate tectonics is too weak to provide the force needed to build the Andes of South America.

North America - a similar story

The mountains of western North America resemble other mountains formed by collision such as the Himalayas. The Rocky Mountains are considered to have formed during the Laramide orogeny from 80 to 45 million years ago, in the latter half of the 180 million year long separation of North America from Africa. Yet western North America "lacks a 'collider'",²⁵ something to push against to cause crustal shortening. Uplift is generally thought to have resulted from friction with the Farallon plate.

Like South America, there is no subducting slab attached to the North American plate to pull on it. The Farallon plate/slab is an eastward extension of the Pacific plate, and now is mostly beneath North America. It is unusual because of the very shallow descent angle attributed to it. In a study of best-fit models, researchers found that the Farallon slab, which extends from the west coast of North America to the east under the continent, dropped passively and was overridden about 1500 km by the North American plate. This indicates "that beneath North America no major drift of the lower mantle has occurred."¹⁷




Farallon plate remnants in cutaway

"Our modelling suggests that North America is strongly coupled to Earth's interior only at the cratonic root [in the large Canadian Shield around and under Hudson Bay], and the NE direction of root drag implies that mantle beneath North America moves at a relatively low speed compared to the SW motion of North America." For the best-fit models, cratonic root basal traction is 40-50 bar going NE-NNE, affecting about 5% of the North American plate area. This "tends to counteract ridge push." "Ridge push is the single most important load acting on the North American plate."¹⁷



Another study estimates the ridge-push force on North America at about 200 bars, and that the total basal forces "are of the same order as ridge push." They believe the basal force is driving the plate forward, not resisting.⁴ While the direction of basal flow has long been disputed,^{5,6,23,35} even the best-case scenario (200 bars ridge push + 200 bars basal drive = 400 bars) is obviously inadequate to raise the vast mountain ranges of western North America.



Tibetan Plateau - where's the driving force?

"The Tibetan Plateau, which formed as a result of the collision between India and Eurasia, has the largest gravitational potential energy (GPE) signal on Earth." "There is no apparent downgoing slab attached to the Indian plate that might assist in driving the plate into Eurasia through the slab pull mechanism. Because the plate is surrounded along its entire southern margin by mid-oceanic ridges, the motion of the Indian plate has been attributed to the ridge push force." "However, the ridge push... is not sufficient."¹³ A 3-dimensional compensated model studied the forces in the region. "The magnitude of stresses associated with GPE differences between Tibet and low-elevation regions in our compensated model is $\sim 2.5 \times 10^{12}$ N/m, while the mid-oceanic ridges exert a force of only $\sim 1 \times 10^{12}$ N/m."¹³ These force per unit-distance numbers show that the force from ridge push needed to just hold up the Tibetan Plateau, not to raise it in the first place, is about 2.5 times too small. "It is clear that something is missing as a driving force that does not have its source within the lithospheric shell."¹³

Other studies find GPE differences between the Tibetan Plateau and surrounding lowlands at 6 to 7×10^{12} N/m¹³ or 7.5 to 8×10^{12} N/m¹⁸, and ridge push at 2.4×10^{12} N/m³⁹. Results from one study determined that around the Tibetan Plateau, a large portion of the lithosphere's strength is in the upper crust, and that a force as high as 1000 to 3000 bars was needed to deform it.¹¹

Let's look at the situation more closely. "Almost all earthquakes on the continents are confined within a crustal layer that varies in thickness from about 10 to 40 km, and are not in the mantle." "The mantle part of the continental lithosphere is relatively weak." "Thus the strength of the continental lithosphere is likely to be contained within the seismogenic layer, variations in the thickness of this strong layer determining the heights of the mountain ranges it can support."²⁴

A team of researchers wrote in 2000, "for almost 20 years the popular view of continental strength

profiles has consisted of a weak lower crust sandwiched between relatively strong layers in the upper crust and mantle. We now believe this view to be incorrect. Earthquake focal depths and gravity anomalies instead suggest that the strength of the continents resides in the seismogenic layer within the crust, and that the continental mantle lithosphere is relatively weak."²⁴

"Moderate-sized earthquakes showing thrust faulting on gently dipping planes in the Himalaya occur at depths of about 15 km, and apparently not deeper than about 18 km." "Earthquakes in continental crust seem to be confined to depths where the temperature is less than 350 to 450 degrees C." "If moderate earthquakes occur at depths no greater than 18 km in the Himalaya because temperatures exceed 350 degrees C along the fault at that depth, then shear stresses of nearly 100 MPa [1000 bars] are necessary to raise the temperature to such a value at that depth."²⁹

Also, to get enough heat for melting the granitoid magmas found near the Main Central Thrust of the Himalayas, "calculations of temperatures appropriate for the Himalaya suggest that shear stresses of 100 MPa [1000 bars] on the Main Central Thrust probably are required to account for the Tertiary granites of the region, if melting took place after slip began on the thrust."²⁹

"The forces driving mountain building must do work against two opposing forces. One might be called mechanical strength, for the stronger the material, the greater the stress needed to deform it." "The forces driving continental plates or blocks together to form mountain ranges must also do work against gravity."³⁰

"Relatively little is known about the absolute strength of the lithosphere and how such strength is distributed with depth." "The oceanic ridges are features with an excess potential energy... of about 1.2×10^{12} N/m." "Estimates of the upper-crustal strength inferred from stress measurements in the KTB (Continental Deepbore Drilling Program) wellhole in Germany show that the cumulative force needed to deform crustal material is in the range of 2 to 5×10^{12} N/m."⁸ If the ridge push force equates to about 200 bars, then crustal deformation alone requires from 330 to 830 bars.

The formation of both mountain ranges and crustal roots creates gravitational potential energy. Moreover, part of this energy increases as the square of the mean elevation, or of the excess thickness of crust in the root. Consequently, as mountain ranges and high plateaus are elevated and the crust is thickened, an increasingly larger amount of work must be done for each subsequent increment of uplift."³⁰

"Because of the increasing amount of work that must be done against gravity acting on an increasingly higher [mountain] range, the range should reach a maximum mean elevation related to the force at which the plates are pushed together. In this sense, the mean elevations of high plateaus serve as crude pressure gauges for the average compressive stresses pushing on the margins of the plateaus. When the maximum elevations are reached, crustal shortening need not cease; convergence can continue as the range builds outward, growing laterally into a high plateau."³⁰

These investigators calculate the required horizontal force to maintain the current elevations to be 690 bars x 100 km depth for Tibet, and 520 bars x 100 km depth for the Andes. "By writing in these units of bars x kilometers, we can visualize these forces per unit length as average horizontal compressive stresses applied to layers of a given thickness. Clearly, changing the value of this thickness requires a

proportional change in the average horizontal stress."³⁰ So confining the stress to the crust, as seems appropriate, changes the force to 1725 bars x 40 km depth for Tibet, and 1300 bars x 40 km depth for the Andes. Using this same estimate, another researcher determined that the "force needed to support Tibet" is "equivalent to average deviatoric stresses of ~120 MPa [1200 bars] if contained within the 40 km elastic layer".²⁴

In 1982, several investigators contemplated the force needed to hold up the Himalayas, using atmospheres (atm) as a measure of pressure. An atm and a bar are almost equivalent. "We considered a model in which the only driving forces are the pulling force of a subducting plate and the pushing force of an oceanic ridge. In that model the stress... operating in a collision boundary was estimated being of the order of 100 atm. However, the Himalayas cannot be held up by such a small stress." The calculated "a value of over 600 atm". "Note, however, that this estimate of 600 atm represents the minimum necessary amount to hold up the Himalayas and Tibet, and is derived by considering these as if they were a tank of water. Thus, within an actual solid body there will probably be much larger stresses."²⁷

As the above examples illustrate, the force needed to deform the crust and raise the Himalayas far exceed the capability of plate tectonics mechanisms.

Slab pull

The slab pull force deserves special consideration because it is central to plate tectonics theory. Here are excerpts from a 50 page review of slab pull and subduction.

"Most of the literature indicates that the slab pull is about 3.3×10^{13} N/m. This is a force per unit-length parallel to the trench."⁹ It is probably not enough force to compress lithosphere and raise mountains.⁹ [paraphrased from original text, p. 156]

"The subduction process is more a passive feature rather than being a driving mechanism of plate motions." Most dipping slabs are being compressed at depths typically shallower than 300 km, indicating slabs forced to sink rather than falling away.⁹


"Some plates... move without any slab pulling them, for example the westward movements of North America, Africa, and South America. Trench suction has been proposed to explain these movements, but beneath both North and South America the mantle is moving relatively eastward, opposite to the kinematics required by the trench suction model." "The African plate moves westward without any slab in its western side. Moreover, it moves opposite to a hypothetical Atlantic ridge push."⁹

"The energy required to pull the plates is far higher than the strength that plates can withstand under extension." The subducted slab is likely able to sustain the load induced by slab pull, but this load probably cannot be transmitted to the rest of the plate without breaking it apart.⁹

"The slab pull concept is based on the hypothesis of a homogeneous composition of the upper mantle, with the lithosphere sinking only because it is cooler. However, the oceanic lithosphere is frozen shallow asthenosphere, previously depleted beneath a mid-oceanic ridge. Depleted asthenosphere is lighter than the 'normal' deeper undepleted asthenosphere. Therefore the assumption that the lithosphere is heavier

only because it is cooler might not be entirely true, and the slab pull could be overestimated."⁹

"Within a slab, eclogitization [(transforming to eclogite under increased pressure and temperature)] is assumed to make the lithosphere denser. However, eclogitization is concentrated in the 6-8 km thick oceanic crust, whereas the remaining 60-80 km-thick lithospheric mantle does not undergo the same transformation. Therefore only 1/10 of the slab is apparently increasing in density, but the main mass of the slab (90%) does not change significantly." "Nevertheless, this type of metamorphic transition is often assumed to be able to determine the slab pull." The small density contrast between subducting crust and mantle casts doubt on the effectiveness of slab pull.⁹



Tomographic images of earth's interior are often used to show subduction. However, "low velocity volumes of the mantle detected by tomography can be due to lateral variations in composition rather than in temperature, i.e., they can be even higher density areas rather than hotter, lighter buoyant material as so far interpreted." "As extreme examples, gold or lead have high density but low seismic velocity. Therefore the interpretation of tomographic images of the mantle where the red (lower velocity) areas are assumed as lighter and hotter rocks can simply be wrong, i.e., they may even be cooler and denser. With the same reasoning, blue (higher velocity) areas, which are assumed as denser and cooler rocks may even be warmer and lighter." Sometimes pieces are assumed to have detached from slabs and fallen away, and appear as blobs on tomographic images. "Tomographic images are based on velocity models that often overestimate the velocity of the asthenosphere where usually the detachment is modeled. Therefore the detachment disappears when using slower velocity for the asthenosphere in

the reference velocity model, or when generating regional tomographic images with better accuracy."⁹

"If oceanic lithosphere is heavier than the underlying mantle, why are there no blobs of lithospheric mantle falling in the upper mantle below the western, older side of the Pacific plate?" Slab dip has been thought to be related to its age, as in "the western Pacific subduction zones because the subducting western Pacific oceanic lithosphere is older, cooler and therefore denser. However, the real dip of the slabs worldwide down to depths of 250 km shows no relation with the age of the downgoing lithosphere."⁹

"There is an observed net westward drift of the lithosphere, with respect to the hotspot reference frame, of 5 to 9 cm per year."³³ This can also be thought of as eastward drift of the mantle. We are supposed to believe that "the 700 km-long West Pacific slab... should pull and carry the 10,000 km-wide Pacific plate, 33 times bigger, overcoming the shear resistance at the plate base and the opposing basal drag induced by the relative eastward mantle flow."⁹

"The negative buoyancy [(dense enough to sink)] of slabs should determine the pull of plates, but it has been shown that the dip of the subduction zones is not correlated with the age and the thermal state of the downgoing plates." "In fact there are slabs where, moving along strike, the age of the downgoing lithosphere varies, but the dip remains the same, or vice versa, the age remains constant while the dip varies (Philippines). There are cases where the age decreases and the dip increases (Western Indonesia), and other subduction zones where the age increases and the dip decreases (Sandwich). This shows that there is not a first order relationship between slab dip and lithospheric age."⁹

Finally, no matter how it is computed, "results do not support a correlation between slab length percentage (length of the trench compared to the length of the whole boundary around the plate) and plate velocity."⁹

Why should the lithosphere start to subduct at all? "Hydrated and serpentized oceanic lithosphere that has not yet been metamorphosed by the subduction process... is still less dense." Slab pull has been calculated to become potentially efficient only at a certain depth, around 180 km. Shallower than that, how is subduction initiated?⁹

"This long list casts doubts on the possibility that slab pull can actually trigger subduction... and drive plate motions."⁹

Another researcher highlights the basic problem for plate tectonics theory

"Why convection in Earth's mantle gives rise to plate tectonics is not obvious. The top thermal boundary layer is supposed to be very stiff because the viscosity of silicate rocks is strongly temperature-dependent. This temperature dependency is so strong that so-called stagnant lid convection should be the most likely mode of mantle convection; the entire surface should be covered by just one single plate, not by a number of rigid plates." "What is needed for the self-consistent generation of plate tectonics is... a mechanism to initiate subduction." Consider moderate-age oceanic lithosphere, say 100 million years old. Even at a "depth range of 10-45 km, oceanic lithosphere of this age is too stiff to be deformed by any reasonable tectonic stress, resulting in a fatal bottleneck for the operation of plate tectonics." "The currently available estimate suggests that yield strength for this 'semi-brittle' regime is still on the order of

600-800 MPa [6000-8000 bars]." "It is known that the self-consistent numerical modeling of plate tectonics (i.e., plate tectonics naturally arising from buoyancy distribution and given rheology, not imposed by boundary conditions) is impossible with such high yield stress. In all previous attempts to simulate plate tectonics in a self-consistent fashion, therefore mantle rheology is modified in one way or another" to get the desired result.²²

1. Alvarez, Walter. August 10, 1982. Geological Evidence For The Geographical Pattern of Mantle Return Flow and the Driving Mechanism of Plate Tectonics. *Journal of Geophysical Research*, Vol. 87, No. B8, pp.6697-6710.
2. Alvarez, Walter. October 1990. Geologic Evidence For The Plate-Driving Mechanism: The Continental Undertow Hypothesis And The Australian-Antarctic Discordance. *Tectonics*, Vol. 9, No. 5, pp.1213-1220.
3. Bird, Peter. May 10, 1998. Testing hypotheses on plate-driving mechanisms with global lithosphere models including topography, thermal structure, and faults. *Journal of Geophysical Research*, Vol. 103, No. B5, pp.10,115-10,129.
4. Bokelmann, Goetz H. R., Paul G. Silver. 2002a. Shear stress at the base of shield lithosphere. *Geophysical Research Letters*, Vol. 29, No. 23, pp. 6-1 to 6-4.
5. Bokelmann, G.H.R. 2002b. Convection-driven motion of the North American craton: Evidence from P-wave anisotropy. *Geophysical Journal International*, Vol. 248, No. 2, pp. 278-287.
6. Bokelmann, G.H.R. November 2002c. Which forces drive North America? *Geology*, Vol. 30, No. 11, pp.1027-1030.
7. Bott, M.H.P. 1993. Modelling the plate-driving mechanism. *Journal of the Geological Society, London*, Vol. 150, pp.941-951.
8. Coblenz, David D., Randall M. Richardson. August 1994. On the gravitational potential of the Earth's lithosphere. *Tectonics*, Vol. 13, No. 4, pp. 929-945.
9. Doglioni, Carlo, Eugenio Carminati, Marco Cuffaro, Davide Scrocca. 2007. Subduction kinematics and dynamic constraints. *Earth Science Reviews*, Vol. 83, pp. 125-175.
10. England, Philip, Gregory Houseman. March 10, 1986. Finite Strain Calculations of Continental Deformation 2. Comparison With the India-Asia Collision Zone. *Journal of Geophysical Research*, Vol. 91, No. B3, pp.3664-3676.
11. Flesch, Lucy M., A. John Haines, William E. Holt. August 10, 2001. Dynamics of the India-Eurasia collision zone. *Journal of Geophysical Research*, Vol. 106, No. B8, pp. 16,435-16,460.
12. Forsyth, Donald, Seiya Uyeda. 1975. On the Relative Importance of the Driving Forces of Plate

Motion. *Geophysical Journal of the Royal Astronomical Society*, Vol. 43, pp.163-200.

13. Ghosh, Attreyee, William E. Holt, Lucy M. Flesch, A. John Haines. May 2006. Gravitational potential energy of the Tibetan Plateau and the forces driving the Indian plate. *Geology*, Vol. 34, No. 5, pp. 321-324.
14. Gung, Yuancheng, Mark Panning, Barbara Romanowicz. 17 April 2003. Global anisotropy and the thickness of continents. *Nature*, Vol. 422, pp.707-711.
15. Guptasarma, D. 2002. Plate Tectonics--Some recent Results, Questions, and Mechanisms. in *Dynamics of Earth's Fluid System*, eds. Rai, S.N., D.V. Ramana, A. Manglik. Lisse.
16. Hamilton, Warren B. 2002. The Closed Upper-Mantle Circulation of Plate Tectonics. in *Plate Boundary Zones, Geodynamics Series Volume 30*, eds. Seth Stein and Jeffrey T. Freymueller, 425 pgs, pp. 359-410.
17. Humphreys, Eugene D., David D. Coblenz. Submitted July 2005. North America Dynamics and Western U.S. Tectonics. Submitted to *Journal of Geophysical Research*, 32 pages.
18. Jimenez-Munt, Ivone, John P. Platt. 2006. Influence of mantle dynamics on the topographic evolution of the Tibetan Plateau: Results from numerical modeling. *Tectonics*, Vol. 25, TC6002, 17 pgs.
19. Kennett, B.L.N. 17 April 2003. Roots of the matter. *Nature*, Vol. 422, pp. 674-675.
20. Kerr, Richard A. 1 September 1995. Earth's Surface May Move Itself...But Did Deeper Forces Act To Uplift the Andes? *Science*, Vol. 269, pp.1214-1216.
21. Kley, Jonas, Tim Vietor. 2007. Subduction and Mountain Building in the Central Andes. in *The seismogenic zone of subduction thrust faults*, eds. Timothy H. Dixon and J. Casey Moore, Columbia University Press, pp. 624-659.
22. Korenaga, Jun. 2007. Thermal cracking and the deep hydration of oceanic lithosphere: A key to the generation of plate tectonics? *Journal of Geophysical Research*, Vol. 112, B05408, 20 pages.
23. Liu, Zhen, Peter Bird. 2002. North America plate is driven westward by lower mantle flow. *Geophysical Research Letters*, Vol. 29, No. 24, pp. 17-1 to 17-4.
24. Maggi, A., J.A. Jackson, D. McKenzie, K. Priestley. June 2000. Earthquake focal depths, effective elastic thickness, and the strength of the continental lithosphere. *Geology*, Vol. 28, No. 6, pp. 495-498.
25. Maxson, Julie, Basil Tikoff. November 1996. Hit-and-run collision model for the Laramide orogeny, western United States. *Geology*, Vol. 24, No. 11, pp. 968-972.
26. Meijer, P.Th., M.J.R. Wortel. July 30, 1992. The Dynamics of Motion of the South American Plate. *Journal of Geophysical Research*, Vol.97, No. B8, pp.11,915-11,931.

27. Miyashiro, Akiho, Keiiti, Aki, A.M.C. Sengor. 1982. *Orogeny*. John Wiley & Sons, New York. Pages 204-5.
28. Molnar, Peter. 8 September 1988. Continental tectonics in the aftermath of plate tectonics. *Nature*, Vol. 335, pp. 131-137.
29. Molnar, Peter, Philip England. April 10, 1990. Temperatures, Heat Flux, and Frictional Stress Near Major Thrust Faults. *Journal of Geophysical Research*, Vol. 95, No. B4, pp. 4833-4856.
30. Molnar, Peter, Helene Lyon-Caen. 1988. Some simple physical aspects of the support, structure, and evolution of mountain belts. *Geological Society of America Special Paper* 218, pp.179-207.
31. Price, Neville J. 2001. *Major Impacts and Plate Tectonics - A model for the Phanerozoic evolution of the Earth's lithosphere*. London.
32. Russo, R.M., P.G. Silver. June 1996. Cordillera formation, mantle dynamics, and the Wilson cycle. *Geology*, Vol. 24, No. 6, pp.511-514.
33. Schellart, W.P. 2004. Quantifying the net slab pull force as a driving mechanism for plate tectonics. *Geophysical Research Letters*, Vol. 31, L07611, 5 pgs.
34. Scoppola, B., D. Boccaletti, M. Bevis, E. Carminati, C. Doglioni. January/February 2006. The westward drift of the lithosphere: A rotational drag? *GSA Bulletin*, Vol. 118, No. 1/2, pp. 199-209.
35. Silver, P.G., W.E. Holt. 8 February 2002. The Mantle Flow Field Beneath Western North America. *Science*, Vol. 295, pp. 1054-1057.
36. Silver, P.G., Raymond M. Russo, Carolina Lithgow-Bertelloni. 2 January 1998. Coupling of South American and African Plate Motion and Plate Deformation. *Science*, Vol. 279, pp.60-63.
37. Smith, Alan D., Charles Lewis. 1999. The planet beyond the plume hypothesis. *Earth-Science Reviews*, Vol. 48, pp.135-182.
38. Smith, Alan D., Charles Lewis. 1999. Differential rotation of lithosphere and mantle and the driving forces of plate tectonics. *Geodynamics*, Vol. 28, pp.97-116.
39. Whittaker, A., M.H.P. Bott, G.D. Waghorn. July 30, 1992. Stresses and Plate Boundary Forces Associated With Subduction Plate Margins. *Journal of Geophysical Research*, Vol. 97, No. B8, pp. 11,933-11,944.
40. Wilson, M. 1993. Plate-moving mechanisms: constraints and controversies. *Journal of the Geological Society*, London, Vol. 150, pp.923-926.

by John Michael Fischer, 2003 - 2008)
www.newgeology.us

